arXiv:1205.0191v1 [math.DS] 1 May 2012

THE ω-LIMIT SETS OF QUADRATIC JULIA SETS

ANDREW D. BARWELL AND BRIAN E. RAINES

Abstract. In this paper we characterize ω-limit sets of dendritic Julia sets for quadratic maps. We use Baldwin's symbolic representation of these spaces as a non-Hausdorff itinerary space and prove that quadratic maps with dendritic Julia sets have shadowing, and also that for all such maps, a closed invariant set is an ω-limit set of a point if, and only if, it is internally chain transitive.

1. Introduction

In this paper we consider the dynamics of quadratic maps on the complex plane, $f_c(z) = z^2 + c$. The interesting dynamics of such a map are carried on the Julia set which is often a strange self-similar space. There are many values of c for which the Julia set is a *dendrite*, a locally connected and uniquely arcwise connected compact metric space. One example is when the value c is strictly pre-periodic; such values are known as Misiurewicz points and the corresponding Misiurewicz maps are well studied (see for example [1, 9, 11]).

Baldwin gives an efficient encoding of the dynamics of these quadratic maps restricted to the Julia set, into a shift map on a non-Hausdorff itinerary space [2]. A detailed description of all such itinerary spaces is given in [4], and an illustration of the utility of non-Hausdorff itinerary spaces in analyzing dynamical systems is given in [3]. Baldwin's collection of itinerary spaces is a large family of dendrites and maps that includes all of the dendritic Julia sets for quadratic maps. It also includes maps on self-similar dendrites which are not Julia sets. Our ω-limit set characterization theorem applies to both the Julia set dendrites and the dendrite maps which are not Julia sets.

A set Λ is *internally chain transitive (ICT)* provided for every $\varepsilon > 0$ and every pair $x, y \in \Lambda$ there is an ε *pseudo-orbit* from x to y in Λ; in other words a collection $\{x = x_0, x_1, \ldots x_n = y\} \subseteq \Lambda$ such that $d(f(x_i), x_{i+1}) < \varepsilon$ for $0 \leq i \leq n-1$. In the family of compact metric spaces this property is equivalent with Sarkovskii's property of weak incompressibility, [8]. We say a set M is *weakly incompressible* provided for every nonempty closed proper subset K of M we have

$$K \cap \overline{f(M-K)} \neq \emptyset.$$

Sarkovskii proved that every ω-limit set has weak incompressibility, and so every ω-limit set is ICT (a fact also proved in [10]). In this paper we consider the question of the converse: *'Is every ICT set an ω-limit set?'*

Let $f : X \to X$ on a metric space (X, d) be continuous. A δ *pseudo-orbit* is a finite or infinite ordered set $\{x_i\}$ such that $d(f(x_i), x_{i+1}) < \delta$ for all i. We say the map f has *shadowing* if for every $\varepsilon > 0$ there is a $\delta > 0$ such that for every δ pseudo-orbit $\{x_i\}$ there is a point $z \in X$ for which $d(f^i(z), x_i) < \varepsilon$ for all i; we say that z ε-*shadows* the pseudo-orbit. In [7] the authors conjecture that the answer to the question *'Is every ICT set an ω-limit set?'* is 'yes' for any map which has shadowing. The conjecture is supported by a result that shows for every tent map with a periodic critical point (maps which have shadowing) the answer is 'yes', and by an example showing there are tent maps without shadowing for which the answer is 'no'. Furthermore, the authors prove in [5] that the answer is 'yes' in the class of shifts of finite type (SFTs), a class of map which has shadowing [12], but the answer is 'no' in the class of sofic shifts (which do not have shadowing in general). There are maps of the unit interval for which the answer is 'yes' and others for which the answer is 'no', but for which shadowing is neither proved nor disproved [6]. Through the use of Baldwin's encoding, we demonstrate that the answer is 'yes' for the class of dendritic Julia sets of quadratic maps, and that these maps have shadowing, further supporting the above conjecture. In fact we prove these results for all self-similar dendrite maps with the unique itinerary property – a large class of maps containing the dendritic Julia set maps.

The paper is organized as follows. In the next section we present the preliminary definitions and results for dendrite maps. In Section Three we prove lemmas regarding the structure and properties of pseudo-orbits in these spaces, and in Section Four we prove shadowing for dendritic Julia sets (Theorem 4.2), from which we obtain the proof of the main theorem, which states that for dendritic Julia sets, every ICT set is necessarily an ω-limit set (Theorem 4.3).

2. Preliminaries

We examine the dynamics of quadratic Julia sets that are dendrites via Baldwin's encoding using non-Hausdorff itinerary spaces. In this section we give a brief description of the definitions and results from

[2] that relate to dendritic Julia sets. More detail and results can be found also in [4].

Given a finite set of symbols, A, for each $n \in \mathbb{N}$ we denote the words of length n from A by A^n, and we define the set of finite words from A by

$$A^{<\omega} = \bigcup_{n \in \mathbb{N}} A^n.$$

For a finite word $\alpha \in A^{<\omega}$ let $len(\alpha) \in \mathbb{N}$ be the length of α.

It is a standard practice in dynamics to encode complicated behavior via symbolic dynamics. A typical simple encoding is to assign symbols from an alphabet to disjoint regions and then to track orbits by recording the regions the points traverse. A difficulty arises due to the fact that the sequence space with its natural topology is totally disconnected while the dynamical system under consideration is usually not. Therefore the encoding usually induces a semiconjugacy rather than a conjugacy. Baldwin avoids this difficulty by the use of "wildcard" symbols. His encoding of the symbol space includes a $*$ symbol which stands for all of the other symbols at once. The topology then on his set of symbols is not the discrete topology, but rather a slightly more complicated non-Hausdorff topology.

Consider the set $\{0, 1, *\}$ with the non-Hausdorff topology with basis $\{\{0\}, \{1\}, \{0, 1, *\}\}$. Let Λ be the product space induced by this topology on $\{0, 1, *\}^\omega$ (ω is the set of non-negative integers, $\{0\} \cup \mathbb{N}$.) For each $\alpha \in \{0, 1, *\}^{<\omega}$, we can define the basic open cylinder sets as

$$B^\Lambda_\alpha \;:=\; \big\{\beta \in \Lambda \;:\; \beta_i = \alpha_i \text{ whenever } \alpha_i \neq *, \text{ for } i \leq len(\alpha)\big\}.$$

Notice that with these definitions, in the factor spaces $*$ cannot be separated via open sets from 1 and 0, so in the product space we cannot separate different infinite words if they only differ in places where one of them has a $*$. This new topology on Λ gives it many connected shift-invariant subspaces. In fact Baldwin showed that it contains copies of every dendritic Julia set for a quadratic map. We make this more precise below.

Let σ denote the natural shift map on a product space. A sequence $\tau \in \Lambda$ is called Λ-*acceptable* if, and only if, the following hold.

1. For all $n \in \omega$, $\tau_n = *$ if, and only if $\sigma^{n+1}(\tau) = \tau$.
2. For all $n \in \omega$ such that $\sigma^n(\tau) \neq \tau$ there is an $m \in \omega$ such that $* \neq \tau_{m+n} \neq \tau_m \neq *$, that is to say if $\sigma^n(\tau) \neq \tau$ then these two sequences differ in a position where neither is $*$.

The sequences τ which are Λ-acceptable are the sequences which we can view as *kneading sequences.* If τ is Λ-acceptable, then $\alpha \in \Lambda$ will be called (Λ, τ)-*consistent* if, and only if, for all $n \in \omega$, $\alpha_n = *$ implies that $\sigma^{n+1}(\alpha) = \tau$. A sequence $\alpha \in \Lambda$ is called (Λ, τ)-*admissible* if, and only if, it is (Λ, τ)-consistent and for all $n \in \omega$ such that $\sigma^n(\alpha) \neq *\tau$ then there is a position where these sequences differ and neither is a star (i.e. there is some $m > 0$ such that $* \neq \alpha_{m+n} \neq \tau_{m-1} \neq *$).

Definition 2.1. Let $\mathcal{D}_\tau := \big\{x \in \Lambda \ : \ x \text{ is } (\Lambda, \tau)\text{-admissible}\big\}$.

Thus $\mathcal{D}_\tau$ is the set of all possible itineraries allowed by the kneading sequence τ, about which Baldwin proved the following:

Theorem 2.2. [2, Theorem 2.4] *Let τ be Λ-acceptable. Then $\mathcal{D}_\tau$ is a shift-invariant self-similar dendrite.*

Baldwin proved that $\mathcal{D}_\tau$ has a natural arc-length (or taxicab) metric, d, which we use in this paper. Moreover, this family of spaces includes all of the quadratic Julia sets which are dendrites:

Theorem 2.3. [2, Theorem 2.5] *Let $f_c(z) = z^2 + c$. If the Julia set of $f_c(z)$, J_c, is a dendrite then there is a Λ-acceptable sequence τ such that $f_c|_{J_c}$ is conjugate to $\sigma|_{\mathcal{D}_\tau}$.*

Let τ be a Λ-acceptable sequence. We call τ the *kneading sequence* for the self-similar dendrite $\mathcal{D}_\tau$. We call the point in $\mathcal{D}_\tau$ of the form $*\tau$ the *critical point* and all of its pre-images *precritical points.*

Let $x = x_0x_1\dots$ and $y = y_0y_1\dots$ be non-precritical points, with $z = z_0z_1\dots$ another (possibly precritical) point, and let $n \in \omega$. We define

$$x{\restriction_n} := x_0x_1\dots x_n.$$

For every $i \leq n$, if $x_i = z_i$ whenever $z_i \neq *$, we say that $x{\restriction_n}$ and $z{\restriction_n}$ are *equivalent*, and we write

$$x{\restriction_n} \approx z{\restriction_n}\ .$$

If instead $x_i = z_i$ for all $i \in \omega$ whenever $z_i \neq *$ then we write

$$x \approx z.$$

We say that

$$x{\restriction_n} \simeq y{\restriction_n}$$

if $x{\restriction_n} = y{\restriction_n}$ or there is a pre-critical point z' for which

$$x{\restriction_n} \approx z'{\restriction_n} \approx y{\restriction_n}\ .$$

Given $\alpha \in \{0, 1, *\}^{<\omega}$, the basic open cylinder sets for $\mathcal{D}_\tau$ are

$$B^\tau_\alpha := \big\{x \in \mathcal{D}_\tau \ : \ x_i = \alpha_i \text{ whenever } \alpha_i \neq *, \text{ for } i \leq len(\alpha)\big\}.$$

Lemmas 2.4 and 2.7 and Remark 2.6 are easy observations.

Lemma 2.4. *Suppose that* z_1 *and* z_2 *are precritical points, and* x *is not precritical with*

$$z_1{\restriction_n}\approx x{\restriction_n}\approx z_2{\restriction_n} .$$

for $n \in \omega$. *Then* $z_1{\restriction_n}\approx z_2{\restriction_n}$.

Definition 2.5. If τ is periodic, let P be the period of τ. If τ is not periodic then for each integer $m \in \mathbb{N}$ we define the *return time for* m, r_m, as

$$r_m = \min\left(\{k \in \mathbb{N} : \sigma^k(\tau){\restriction_m}= \tau{\restriction_m}\} \cup \{\infty\}\right).$$

It is easy to see that if τ is non-recurrent then there is some M such that for all $m \geq M$, $r_m = \infty$. Also if τ is recurrent but not periodic then r_m is an integer for each $m \in \mathbb{N}$ and $r_m \to \infty$ as $m \to \infty$.

Remark 2.6. Notice that if $k, j \in \mathbb{N}$, with $k > j$ such that

$$\tau{\restriction_j}\approx \sigma^{k-j}(\tau){\restriction_j},$$

then

1. if τ is periodic with period P we either have that $k - j = nP$ for some $n \in \mathbb{N}$, or $j < P$;
2. if τ is not periodic we have that $k - j \geq r_j$.

The next result is central to our understanding of dendrite maps, and follows from the definition of the cylinder sets above.

Lemma 2.7. *For* $x, y \in \mathcal{D}_\tau$ *and* $\varepsilon > 0$, *there is an* $N_\varepsilon \in \mathbb{N}$ *such that*

$$d(x, y) < \varepsilon$$

if, and only if

$$x{\restriction_{N_\varepsilon}}\simeq y{\restriction_{N_\varepsilon}} .$$

This lemma describes the key difference between the setting of this paper and the setting of shifts of finite type. In SFTs two points are within ε of each other if and only if they are *equal* for their initial sement of length N_ε. In the symbolic spaces we study in this paper, we know that two points x and y are within ε of each other if, and only if either:

1. x and y have exactly the same initial segment of length N_ε, or
2. x and y have a disagreement in say the jth place (with $j \leq N_\varepsilon$), in which case there is a precritical point z between them that has a $*$ in the jth place.

Definition 2.8. In case (2) we say that x and y have a *flip* in the jth place.

For example we could have two points x and y within ε of each other as below:

$$\begin{aligned} x &= x_0x_1x_2\dots x_{j-1}x_jx_{j+1}\dots x_{N_\varepsilon} \\ z &= x_0x_1x_2\dots x_{j-1} * \tau_1\tau_2\dots\tau_{N_\varepsilon-j} \\ y &= x_0x_1x_2\dots x_{j-1}y_jy_{j+1}\dots y_{N_\varepsilon} \end{aligned}$$

where j is chosen minimally. In this case we must have that the words $x_{j+1}\dots x_{N_\varepsilon}$ and $y_{j+1}\dots y_{N_\varepsilon}$ are equivalent to $\tau_1\dots\tau_{N_\varepsilon-j}$. So if τ is not periodic then it must be the case that these three words are equal. If instead τ is periodic, say with period P, then these words must be equal for the first P many symbols, then they can have another flip, then they must be equal for the next P many symbols, etc. Analyzing this situation for δ pseudo-orbits and using it to construct ε-shadowing points is the focus of the next section.

3. Pseudo-Orbits in Dendrites

In this section we will prove a number of results pertaining to pseudo-orbits in dendrites, which will ultimately allow us to prove that dendrite maps have shadowing.

Let $\delta > 0$. For the purpose of illustration, we begin with a consideration of δ pseudo-orbits in SFT spaces. Let$\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit in X, a SFT. Let $N_\delta \in \mathbb{N}$ be defined so that

$$d(x,y) < \delta$$

if, and only if

$$x\restriction_{N_\delta} = y\restriction_{N_\delta}\ .$$

By definition we have

$$\sigma(x_1)\restriction_{N_\delta} = x_2\restriction_{N_\delta}$$

and

$$\sigma(x_2)\restriction_{N_\delta} = x_3\restriction_{N_\delta}$$

which implies

$$\sigma^2(x_1)\restriction_{n_\delta-1} = \sigma(x_2)\restriction_{N_\delta-1} = x_3\restriction_{N_\delta-1}\ .$$

If we denote x_i by $x^i_0x^i_1x^i_2\dots$ then we have the following array of initial segments of length N_δ where down each column we have equality:

$$\begin{array}{l}
x_0^1x_1^1x_2^1x_3^1x_4^1\dots x_{N_\delta}^1 \\
\quad x_0^2x_1^2x_2^2x_3^2x_4^2\dots x_{N_\delta}^2 \\
\qquad x_0^3x_1^3x_2^3x_3^3x_4^3\dots x_{N_\delta}^3 \\
\qquad\quad x_0^4x_1^4x_2^4x_3^4x_4^4\dots x_{N_\delta}^4 \\
\qquad\qquad x_0^5x_1^5x_2^5x_3^5x_4^5\dots x_{N_\delta}^5
\end{array}$$

The column equality in the above array allows a straightforward proof of shadowing in SFT spaces. For sufficiently small δ the point $z = x_0^1x_0^2x_0^3\dots$ is in the space, and it also δ shadows the pseudo-orbit. It is easy to see from the array that

$$z\restriction_{N_\delta} = x_1\restriction_{N_\delta} .$$

This is also true for the jth shift of z and the point x_j.

The symbolic spaces we consider in this paper are far less simple. Let $\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit in $\mathcal{D}_\tau$, and let $N_\delta \in \mathbb{N}$ be given as in Lemma 2.7. Then, as described after the lemma, for each $i \in \mathbb{N}$ we have

$$\sigma(x_i)\restriction_{N_\delta} \simeq x_{i+1}\restriction_{N_\delta} .$$

This implies that either these two initial segments are equal or there is a flip in some position, and a precritical point, z, between $\sigma(x_i)$ and x_{i+1}.

Consider now the array associated with the δ pseudo-orbit:

$$\begin{array}{l}
x_0^1x_1^1x_2^1x_3^1x_4^1\dots x_{N_\delta}^1 \\
\quad x_0^2x_1^2x_2^2x_3^2x_4^2\dots x_{N_\delta}^2 \\
\qquad x_0^3x_1^3x_2^3x_3^3x_4^3\dots x_{N_\delta}^3 \\
\qquad\quad x_0^4x_1^4x_2^4x_3^4x_4^4\dots x_{N_\delta}^4 \\
\qquad\qquad x_0^5x_1^5x_2^5x_3^5x_4^5\dots x_{N_\delta}^5 \\
\qquad\qquad\quad \ddots\ \ddots\ \ddots\qquad \ddots
\end{array}$$

Since this is a δ pseudo-orbit we know that

$$\sigma^t(x_1)\restriction_{N_\delta-t} \approx \sigma^{t-1}(x_2)\restriction_{N_\delta-t} \approx \dots \approx x_{t+1}\restriction_{N_{\delta-t}} \tag{1}$$

for every $0 \le t \le N_\delta$. Notice also that

$$\sigma^t(x_1)\restriction_{N_\delta-t} = \sigma^{t-1}(x_2)\restriction_{N_\delta-t} = \dots = x_{t+1}\restriction_{N_{\delta-t}} \tag{2}$$

for all $t \leq N_\delta$ if, and only if, we have column equality in these segments as in the SFT case. But by Lemma 2.7 it may be the case that there is a flip in some column, say the jth column, relative to x_1. If j is minimal in this respect, then for all $1 \leq \ell < j$ we have the following column equality in the array:

$$x^1_\ell = x^2_{\ell-1} = \cdots = x^{\ell+1}_0.$$

This does not hold in the jth column, since for some $1 \leq i \leq j$ we have

$$x^1_j = x^2_{j-1} = \cdots = x^i_{j-i+1} \neq x^{i+1}_{j-i}.$$

This motivates the following definition:

Definition 3.1. *We will refer to any such j above as a flip column relative to x_1 and i as a flip row relative to x_1. Given x_t for $t \geq 1$ we define flip columns relative to x_t in an analogous manner, and given a flip column j relative to x_t, we define the flip row of the flip column j to be the least i such that*

$$x^t_j \neq x^{t+i}_{j-i}.$$

These flip rows and flip columns are the main obstruction to proving shadowing and characterizing ω-limit sets for the dendrite maps under consideration.

In SFTs, letting $\delta = \varepsilon$, one can prove every δ pseudo-orbit is ε shadowed by a point like z defined above [12]. We would like to mirror the proof from SFT spaces in dendrites, but we must make some significant changes to account for the flip columns. Given $\varepsilon > 0$ we will choose a $\delta > 0$ (based on several upcoming lemmas) much smaller than ε. Then, given a δ pseudo-orbit $\{x_i\}_{i\in\mathbb{N}}$, we will construct a "canonical form" of an ε-shadowing point, $z = z_0 z_1 \ldots$ where $z_0 = x^1_0$ and if t_1 is the first flip column for the pseudo-orbit we let $z_j = x^{j+1}_0$ for all $j < t_1$ but we define $z_{t_1} = \diamond$. Letting t_2 be the next flip column we define $z_k = x^{k+1}_0$ for all $t_1 < k < t_2$ and again we assign $\diamond$ to z_{t_2}. Continuing this procedure gives us a point $\hat{z}$ with possibly infinitely many places where there is a $\diamond$, but since $\diamond$ is not in our alphabet, $\hat{z}$ is not a point in $\mathcal{D}_\tau$. We must then assign either a 0, 1 or $*$ to each $\diamond$ in $\hat{z}$ in such a way that the resulting point z is both in $\mathcal{D}_\tau$ and ε-shadows the pseudo-orbit.

It is the goal of the next several results to prove that there is a δ small enough so that the $\diamond$'s will occur with larger than N_ε gaps between them. Once this has been established, we then prove that *any* assignment of 0's or 1's to the $\diamond$'s in $\hat{z}$ will generate a true ε-shadow z of the δ pseudo-orbit (unless $\sigma^k(\hat{z}) \approx \tau$ for some $k \in \omega$, in which case there is a unique assignment).

To begin, we prove a basic feature of column equality in arrays related to δ pseudo-orbits between successive flip columns. The proof is immediate from the definition of flip column (see Figures 1 and 2).

Lemma 3.2. *Let* $\{x_i\}_{i\in\mathbb{N}}$ *be a* δ *pseudo-orbit, and let* $k \in \mathbb{N}$ *be chosen such that there are two flip columns,* j_1 *and* j_2 *relative to* x_k *with* $j_1 < j_2$ *and no flip columns relative to* x_k *between* j_1 *and* j_2*. Then we have the following equalities:*

(1) *for all* $1 \leq \ell \leq k + j_1$,

$$\sigma^{j_1+1}(x_k)\restriction_{j_2-j_1-1} = \sigma^{j_1+1-\ell}(x_{k+\ell})\restriction_{j_2-j_1-1},$$

and

(2) *for all* $1 \leq n \leq j_2 - j_1 - 1$

$$\sigma^{j_1+n}(x_k)\restriction_{j_2-j_1-n} = x_{k+j_1+n}\restriction_{j_2-j_1-n} .$$



FIGURE 1. Lemma 3.2 (1).

Before proceeding with a proof that these dendrite maps have shadowing, we need to carefully consider both periodic and non-periodic kneading sequences and the gap between successive flip columns. First we consider the periodic case. Let τ be periodic, $\delta > 0$ and let N_δ be defined as in Lemma 2.7. Let $\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit. Since τ is periodic, if x_k has a flip column in say the jth column with $j < N_\delta$, then it can have a flip column in every $j + kP$ column afterwards, but



FIGURE 2. Lemma 3.2 (2).

also in some columns which are not "in sync" with the period. The next lemma addresses the arrangement of the flip columns that are "out of sync".

Lemma 3.3. *Let $\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit, and let k and $j < N_\delta$ be chosen such there there is a pre-critical point z with*

$$\sigma(x_k)\restriction_{N_\delta} \approx z\restriction_{N_\delta} \approx x_{k+1}\restriction_{N_\delta}$$

and j is minimal with $\sigma^j(z) = *\tau_1\tau_2\ldots\tau_{P-1}*\tau_1\ldots.$ Suppose also that there are flip columns t for x_k such that $t-j$ is not a multiple of P.

Choose s_1 minimal such that s_1 is a flip row of a flip column t_1 with $t_1 - j$ not a multiple of P and

$$t_1 < N_\delta.$$

Given s_i and t_i, define $s_{i+1} > s_i$ to be minimal such that s_{i+1} is a flip row of a flip column t_{i+1} with $t_{i+1} - j$ not a multiple of P and with

$$t_{i+1} < t_i.$$

Then there are finitely many such s_i's and t_i's, and for each such i we have

$$N_\delta - t_i < iP.$$

Proof. We start with the flip row s_1 and flip column t_1. Since s_1 is the least flip row associated with x_k with $t_1 - j$ not a multiple of P, there are no flips in the array in non-$j+kP$ places between $\sigma^{s_1}(x_k)$ and x_{k+s_1} until the t_1 symbol. There must be a precritical point z_1 with

$$\sigma(x_{k+s_1-1})\upharpoonright_{N_\delta} \approx z_1\upharpoonright_{N_\delta} \approx x_{k+s_1}\upharpoonright_{N_\delta}$$

with

$$\sigma^{t_1-s_1}(z_1) = \tau.$$

But since t_1 is a flip column for x_k and s_1 is the least flip row, it must be the case that

$$\sigma^{t_1+1}(z) = \sigma^{t_1+1-j}(\tau) \neq \tau_1\tau_2\ldots$$

but

$$\sigma^{t_1+1-j}(\tau)\upharpoonright_{N_\delta-t_1} \approx \sigma^{t_1+1-s_1}(x_{k+s_1})\upharpoonright_{N_\delta-t_1}$$

and

$$\sigma^{t_1+1-s_1}(x_{k+s_1})\upharpoonright_{N_\delta-t_1} \approx \sigma^{t_1+1-s_1}(z_1)\upharpoonright_{N_\delta-t_1} = \tau_1\tau_2\ldots\tau_{N_\delta-t_1}.$$

Therefore

$$\sigma^{t_1+1-j}(\tau)\upharpoonright_{N_\delta-t_1} \approx \tau_1\tau_2\ldots\tau_{N_\delta-t_1}.$$

By Remark 2.6(1) it must be the case that $N_\delta - t_1 < P$.



Figure 3. Lemma 3.3.

The result follows by induction noticing that $\sigma^{t_i-s_i}(x_{k+s_i})$ will agree with $\sigma^{t_i}(x_k)$ on a word of length $t_{i-1}-t_i$ which by similar reasoning must be less than P. We combine this with $N_\delta - t_{i-1} < (i-1)P$ to see that

$$N_\delta - t_i = (N_\delta - t_{i-1}) + (t_{i-1} - t_i) < (i-1)P + P = iP.$$

□

Next we consider the case that $\tau = *\tau_1\tau_2\dots$ is not periodic. The following combinatorial argument will be a useful tool when dealing with return times. For words α and β, when we write $\alpha\beta$ we mean α concatenated with β; for $n \in \mathbb{N}$, when we write α^n we mean α concatenated with itself n times.

Lemma 3.4. *Let α be a word of length n, let $\beta = \alpha\alpha$, and suppose that for some $1 \leq m < n$ we have that $\sigma^{n-m}(\beta)$ begins with the word β. Then there is some word γ of length ℓ with $1 \leq \ell \leq m$, such that ℓ divides m and $n - m$ and $\alpha = \gamma^{n/\ell}$.*

Proof. Set $m_1 = m$ and $\alpha_0 = \alpha$. From the statement of the lemma, we get that $\sigma^{n-m_1}(\beta)$ begins with the initial m_1-segment of α_0; call this segment α_1. Then $\sigma^n(\beta)$, which is also the word α_0, begins with α_1 and this also corresponds to the second m_1-segment of the occurrence of α_0 beginning at $\sigma^{n-m_1}(\beta)$, provided $n \geq 2m_1$ (see Figure 4).



FIGURE 4. Overlapping words.

Continuing in this way we see that either

(1) m_1 divides n, so $\alpha_0 = \alpha_1^{n/m_1}$, or
(2) $\alpha_0 = (\alpha_1^{k_1})\alpha_2$ for some $k_1 \geq 1$ and some word α_2 of length $m_2 < m_1$.

In case (1) we are done. In case (2), if we let $\beta_1 = \alpha_1\alpha_1$, from the structure of the word B we get that $\sigma^{m_1-m_2}(\beta_1)$ begins with α_1. But this is precisely the situation we started with, so we can iterate the above argument. Since every finite word is made up of component symbols, this process cannot continue indefinitely, so there must be some word α_r of length $m_r \geq 1$ such that $\alpha_0 = \alpha_r^{n/m_r}$.

To see that m_r divides m_1, notice that $n = k_1m_1 + m_2$, where $m_2 < m_1$, and similarly $m_1 = k_2m_2 + m_3$ for $k_2, m_3 \in \mathbb{N}$ with $m_3 < m_2$. This is the Euclidean Algorithm, which repeats with m_i being the length of α_i, until we get that $m_{r-1} = k_rm_r$, and m_r is the greatest common divisor of n and m_1. □

Lemma 3.5. *Suppose that τ is not periodic. Then the return times $\{r_m\}_{m\in\mathbb{N}}$ have the property that $r_t \geq t$ for infinitely many $t \in \mathbb{N}$.*

Proof. Recall first that $r_m \leq r_{m+1}$ for every $m \in \mathbb{N}$. Suppose that the statement of the lemma fails; in other words we have that $r_m < m$ for cofinitely many $m \in \mathbb{N}$. Then we can choose m' maximum such that $r_{m'} \geq m'$, and we get that $r_{m'+1} < m'+1$. Thus

$$m' \leq r_{m'} \leq r_{m'+1} \leq m',$$

$$\text{so } r_{m'} = r_{m'+1} = m'.$$

This gives us that the first m'-segment of τ (which follows the initial symbol $*$ of τ) is repeated immediately; let us refer to this segment as α.

Let $i \geq 2$ be minimal such that $r_{m'+i} \neq m'$. Then

$$m' = r_{m'+i-1} < r_{m'+i} < m'+i,$$

$$\text{and so } r_{m'+i} \leq m'+i-1.$$

Let $r_{m'+i} = m'+j$ for $1 \leq j < i$, and pick $k \in \mathbb{N}$ such that $(k-1)m' < m'+j \leq km'$ (see Figure 5).



Figure 5. Return times overlapping.

We claim that for any such i, the $(m'+i)$-segment of τ following the first $(r_{m'+i})$-segment will be an exact repeat of the first $(r_{m'+i})$-segment. This will force the m'-segment after the initial symbol of τ to repeat indefinitely, which is not an acceptable sequence for τ, and the resulting contradiction would complete the proof. We prove that this occurs for the current i under consideration, and a simple argument will conclude that it occurs for all $r_m > r_{m'+i}$.

If $m'+j = km'$ we can stop immediately. Otherwise $(k-1)m' < m'+j < km'$, then Lemma 3.4 tells us that there is some $\ell \leq m'$ such that each identical m'-segment of τ is composed of ℓ identical words of length m'/ℓ. In this case we have that ℓ divides $(k-1)m'-j$ and we have the required repetition.

To conclude the proof, notice that for any $k \geq 1$,

$$r_{m'+i+k-1} \leq r_{m'+i+k} \leq m'+i+k-1,$$

so when the initial $(m'+i+k)$-segment (following the initial symbol) first repeats in τ it does so before the end of the repeat of the first

$(m' + i + k - 1)$-segment. This forces an overlap, and as above we get that the first repeat of the initial $(m' + i + k)$-segment must exactly repeat what has come before it. $\square$

Lemma 3.6. *Let* $\{x_i\}_{i \in \mathbb{N}}$ *be a* δ *pseudo-orbit, let* τ *be a recurrent non-periodic kneading sequence, and let* $\{m_i\}_{i \in \mathbb{N}}$ *be a sequence of natural numbers chosen such that*

$$m_i \leq r_{m_i} < r_{m_i} + 1 < m_{i+1}.$$

Suppose that k *is chosen such that there are two successive flip columns* $j_1 < j_2$ *relative to* x_k *with flip rows* i_1 *and* i_2*. If there exists some* $t \in \mathbb{N}$ *such that*

(i) $j_2 - j_1 - 1 < m_t$, *and*
(ii) $N_\delta - j_2 - 1 > m_{t+1}$,

then there is a third flip column, $j_3 > j_2$*, relative to* x_k *with flip row* i_3 *such that*

(a) i_3 *is between* i_1 *and* i_2*, and*
(b) $j_3 - j_2 - 1 < m_{t+1}$.

Proof. Suppose we have the points, flip columns and rows as described in (i) and (ii) of the lemma, but suppose that there is no third flip column, j_3 satisfying (a) and (b).

Without loss of generality, assume that $i_1 < i_2$. Then there is some pre-critical point $z_1 \in [x_{k+i_i-1}, x_{k+i_1}]$ such that

$$\sigma(x_{k+i_1-1})\restriction_{N_\delta} \approx z_1\restriction_{N_\delta} \approx x_{k+i_1}\restriction_{N_\delta},$$

and $\sigma^{j_1-i_1}(z_1) = \tau = *\tau_1\tau_2\tau_3\ldots$. Hence

$$\sigma^{j_1-i_1+1}(x_{k+i_1})\restriction_{N_\delta-j_1} \approx \tau_1\tau_2\tau_3\ldots\restriction_{N_\delta-j_1}.$$

Similarly, there is a pre-critical point $z_2 \in [x_{k+i_2-1}, x_{k+i_2}]$ such that

$$\sigma(x_{k+i_2-1})\restriction_{N_\delta} \approx z_2\restriction_{N_\delta} \approx x_{k+i_2}\restriction_{N_\delta},$$

and $\sigma^{j_2-i_2}(z_2) = \tau = *\tau_1\tau_2\tau_3\ldots$. Hence

$$\sigma^{j_2-i_2+1}(x_{k+i_2})\restriction_{N_\delta-j_2} \approx \tau_1\tau_2\tau_3\ldots\restriction_{N_\delta-j_2}.$$

We see that

$$\tau_{j_2-j_1+1}\tau_{j_2-j_1+2}\cdots = \sigma^{j_2-j_1+1}(\tau) = \sigma^{j_2-i_1+1}(z_1)$$

and

$$\sigma^{j_2-i_1+1}(z_1)\restriction_{N_\delta-j_2-1} \approx \sigma^{j_2-i_1+1}(x_{k+i_1})\restriction_{N_\delta-j_2-1}.$$

While in the same column we have

$$\tau_1\tau_2\cdots = \sigma^{j_2-i_2+1}(z_2),$$

and

$$\sigma^{j_2-i_2+1}(z_2)\upharpoonright_{N_\delta-j_2-1}\approx \sigma^{j_2-i_2+1}(x_{k+i_2})\upharpoonright_{N_\delta-j_2-1}\ .$$

Since we assume that there are no successive flip columns j_3 with related flip row i_3 where $i_1 < i_3 < i_2$ and we assume $j_3 - j_2 - 1 < m_{t+1}$ and $N_\delta - j_2 - 1 > m_{t+1}$, it must be the case that

$$\begin{aligned}(3)\qquad \sigma^{j_2-i_1+1}(z_1)\upharpoonright_{M}\approx \sigma^{j_2-i_1+1}(x_{k+i_1})\upharpoonright_{M}&\approx\\ \sigma^{j_2-i_2+1}(x_{k+i_2})\upharpoonright_{M}&\approx \sigma^{j_2-i_2+1}(z_2)\upharpoonright_{M}\end{aligned}$$

for some $M \in \mathbb{N}$ with $M \geq m_{t+1}$ (otherwise there *would* have been a flip column j_3 with flip row i_3 satisfying (a) and (b)). There are three cases to consider. For instance we could have

(A) $z_1 \neq x_{k+i_1}$ and x_{k+i_1} is precritical, or
(B) $z_1 = x_{k+i_1}$, or
(C) $z_1 \neq x_{k+i_1}$ and x_{k+i_1} is not precritical.

In case (A), let s be such that $\sigma^{s-i_1}(x_{k+i_1}) = \tau$. Then because j_1 and j_2 are successive flip columns relative to x_k, it must be the case that $s > j_2$. If s is small enough to still be a flip column for x_k then we know that $s - j_2 - 1 \geq m_{t+1}$ by our assumptions that no flip column for x_k satisfies (a) and (b). If s is so large that it cannot be counted as a flip column relative to x_k then it must be the case that $s > N_\delta$. So we again have $s - j_2 - 1 \geq m_{t+1}$. This implies that the first M-length word of $\sigma^{j_2-i_1+1}(z_1)$ equals the first M word of $\sigma^{j_2-i_1+1}(x_{k+i_1})$. So we have

$$\tau_{j_2-j_1+1}\tau_{j_2-j_1+2}\dots\tau_{j_2-j_1+M} = \sigma^{j_2-i_1+1}(x_{k+i_1})\upharpoonright_{M}\ .$$

Cases (B) and (C) lead to the same conclusion via simpler reasoning.

By a similar argument, considering three cases for z_2 and x_{k+i_2}, we see that

$$\tau_1\tau_2\dots\tau_M = \sigma^{j_2-i_2+1}(x_{k+i_2}).$$

So by equation (3) we see that

$$\tau_{j_2-j_1+1}\tau_{j_2-j_1+2}\dots\tau_{j_2-j_1+M} \approx \tau_1\tau_2\dots\tau_M.$$

since τ is non-periodic, none of these symbols can be $*$. Therefore,

$$\tau_{j_2-j_1+1}\tau_{j_2-j_1+2}\dots\tau_{j_2-j_1+M} = \tau_1\tau_2\dots\tau_M.$$

Since $M \geq m_{t+1}$ it then must be the case that $j_2 - j_1 + 1 \geq r_{m+1}$, but we assumed that $j_2 - j_1 - 1 < m_t$ which implies that $j_2 - j_1 + 1 \leq m_t + 1 \leq r_{m_t} + 1 < m_{t+1} \leq r_{m_{t+1}}$, a contradiction. $\square$

We will now define sequences $\{\alpha_k\}_{k\in\mathbb{N}}$, $\{\beta_k\}_{k\in\mathbb{N}}$, $\{j_k\}_{k\in\mathbb{N}}$, and $\{i_k\}_{k\in\mathbb{N}}$, that keep track of the flip columns and rows in a δ pseudo-orbit as follows.



FIGURE 6. Return times in the δ pseudo-orbit of Lemma 3.6; '$\star$' represents a flip.

Definition 3.7. Let $\varepsilon > 0$, and let N_ε be given as in Lemma 2.7. Let $\delta > 0$, and let $\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit, with $x_i = x_0^i x_1^i \ldots$ for each $i \in \mathbb{N}$. Let α_1 be minimal such that there exists a $j_1 < N_\varepsilon$ minimal for which either

(1) there exists $i_1 \leq j_1$, minimal, such that j_1 is a flip column relative to α_1 and i_1 is the flip row of the flip column j_1, in other words
$$x_{j_1}^{\alpha_1} \neq x_{j_1-i_1}^{\alpha_1+i_1};$$
or

(2) we have that
$$x_{j_1}^{\alpha_1} = x_{j_1-i}^{\alpha_1+i} = *$$
for all $1 \leq i \leq j_1$ in which case we define $i_1 = 1$.

Let $\beta_1 = \alpha_1 + j_1$. For $n > 1$ define α_n, j_n, and i_n recursively as above so that $\alpha_n > \beta_{n-1}$. Thus we only start looking for the next point with a flip column in its initial N_ε-segment after we have passed the previous flip column.

These sequences of α_k's and β_k's allow us to keep track of the key flip columns in the δ pseudo-orbit. Specifically, in trying to construct a shadowing point for this δ pseudo-orbit we need to know where the flip columns affect the N_ε agreement. Notice that $\sigma(x_i)\restriction_{N_\delta}\simeq x_{i+1}\restriction_{N_\delta}$, for every $i\in\mathbb{N}$, while our goal is to get a point z which has $\sigma^i(z)\restriction_{N_\varepsilon}\simeq x_i\restriction_{N_\varepsilon}$, for all $i\in\mathbb{N}$, with N_ε much smaller than N_δ. The α_k's give us the points in the pseudo-orbit that have a flip column somewhere in their first N_ε-many symbols, while the β_k's are the actual place where the canonical shadowing point must have a $\diamond$ symbol.

The next result will help us prove that for every $\varepsilon>0$ there is a $\delta_\varepsilon>0$ small enough to guarantee that after the $\diamond$ symbol (in some β_kth place) in our canonical shadowing point, there is a well-defined string of length N_ε that agrees with τ. This means that the actual shadowing point will have its β_kth shift within ε of the critical point. This property will allow us to freely choosing 0 or 1in place of each $\diamond$, as either choice is allowed in these self-similar dendrite maps.

Proposition 3.8. *Let $\varepsilon>0$ and $N_\varepsilon\in\mathbb{N}$ be given. Then there is some $\delta_\varepsilon>0$ such that for any $0<\delta\le\delta_\varepsilon$ and any δ pseudo-orbit, $\{x_i\}_{i\in\mathbb{N}}$ we have*

$$x_{\beta_k+t}\restriction_{N_\varepsilon-t}\approx\sigma^t(*\tau)\restriction_{N_\varepsilon-t}$$

for all $k\in\mathbb{N}$ and all $0\le t\le N_\varepsilon$.

Proof. We prove the proposition in three parts. First we assume that τ is periodic with period P, then we assume that τ is non-recurrent, and finally we assume that τ is recurrent but non-periodic.

To begin, let τ be periodic with period $P\in\mathbb{N}$. So

$$\tau=*\tau_1\tau_2\dots\tau_{P-1}*\tau_1\dots$$

In this case we choose $\delta_\varepsilon>0$ so that

$$N_{\delta_\varepsilon}>2(P+1)N_\varepsilon.$$

Choose $0<\delta\le\delta_\varepsilon$ and $N_\delta\in\mathbb{N}$ via Lemma 2.7. Let $k\in\mathbb{N}$ and consider α_k. First we consider the case that α_k satisfies case (1) of Definition 3.7. Let flip column j_k and associated flip row i_k be as defined. We prove that there are no flips relative to $x_{\alpha_k+i_k}$ between j_k-i_k+1 and $(j_k-i_k+1)+N_\varepsilon$ with flip rows between 1 and $(\beta_k-i_k)+N_\varepsilon$. Lemma 3.2 will imply the proposition.

As in Lemma 3.3, let $1\le s_1\le(\beta_k-i_k)+N_\varepsilon$ be the minimal flip row for a flip column, t_1, relative to $x_{\alpha_k+i_k}$ where t_1-j_k is not a multiple of P. Given s_q and t_q, define $s_q<s_{q+1}\le(\beta_k-i_k)+N_\varepsilon$ to be the least flip row associated with a flip column, t_{q+1}, relative to $x_{\alpha_k+i_k}$ such that $t_{q+1}<t_q$ and such that $t_{q+1}-j_k$ is not a multiple of P. Then

$$N_\delta-t_q<qP$$

for all such q. Let r be the number of flip rows and columns relative to $x_{\alpha_k+i_k}$ enumerated as above. Then r is obviously bounded by the number of rows between i_k and $j_k - N_\varepsilon$. This is maximized when $i_k = 1$ and $j_k = N_\varepsilon - 1$. In that case we have $r \leq N_\varepsilon - 1 + N_\varepsilon < 2N_\varepsilon$. Therefore the last flip column in our enumeration, t_r satisfies

$$N_\delta - t_r < rP < 2N_\varepsilon P.$$

We prove that $t_r > j_k + N_\varepsilon$. Since

$$N_\delta > 2PN_\varepsilon + 2N_\varepsilon,$$

we have

$$t_r > N_\delta - rP \geq N_\delta - 2N_\varepsilon P > 2N_\varepsilon.$$

Since $j_k < N_\varepsilon$,

$$t_r > 2N_\varepsilon > j_k + N_\varepsilon.$$

The case that α_k satisfies (2) of Definition 3.7 is more straightforward. This completes the proof for the periodic case.

Now assume that τ is non-recurrent. Let M be large enough so that $r_m = \infty$ for all $m \geq M$. That is to say the word $\tau_1 \dots \tau_M$ never re-occurs in τ. Let $\delta_\varepsilon > 0$ be chosen so that

$$N_{\delta_\varepsilon} > 2(N_\varepsilon + M + 1).$$

Let $0 < \delta \leq \delta_\varepsilon$ and let $N_\delta \in \mathbb{N}$ be defined as in Lemma 2.7. Let $k \in \mathbb{N}$ and consider α_k. Suppose that α_k satisfies (1) of Definition 3.7. Let j_k and i_k be the flip column and flip row as defined. We prove that there are no flips relative to $x_{\alpha_k+i_k}$ with flip columns between $j_k - i_k + 1$ and $(j_k - i_k + 1) + N_\varepsilon$ and flip rows between 1 and $(\beta_k - i_k) + N_\varepsilon$. Let z be the precritical point that is responsible for the α_k flip. That is to say

$$\sigma(x_{\alpha_k+i_k-1})\restriction_{N_\delta} \approx z\restriction_{N_\delta} \approx x_{\alpha_k+i_k}\restriction_{N_\delta}$$

with

$$\sigma^{j_k-i_k+1}(z) = *\tau_1\tau_2\dots.$$

We have

$$\begin{aligned}\sigma^{j_k-i_k+2}(z)\restriction_{N_\delta-N_\varepsilon-2} &= \tau_1 \dots \tau_{N_\varepsilon+2M} \dots \tau_{N_\delta-N_\varepsilon-2} \\ &= \sigma^{j_k-i_k+2}(x_{\alpha_k+i_k})\restriction_{N_\delta-N_\varepsilon-2}\end{aligned}$$

because $j_k - i_k < N_\varepsilon$. Suppose that there is some other flip relative to $x_{\alpha_k+i_k}$ with flip column between $j_k - i_k + 1$ and $(j_k - i_k + 1) + N_\varepsilon$. Let r be minimal such that the flip row for this column occurs at $x_{\alpha_k+i_k+r}$ and let t be chosen so that the precritical point z' responsible for this flip has

$$\sigma(x_{\alpha_k+i_k+r-1})\restriction_{N_\delta} \approx z'\restriction_{N_\delta} \approx x_{\alpha_k+i_k+r}\restriction_{N_\delta}$$

with

$$\sigma^t(z) = *\tau_1\tau_2\dots$$

and $j_k - i_k - r < t \leq (j_k - i_k - r) + N_\varepsilon$. Since r is the first row with a flip, it must be the case that

$$\begin{aligned}\sigma^{t+1}(z')\restriction_M &= \tau_1 \dots \tau_M \\ &= \sigma^{(j_k-i_k+1)+t-(j_k-i_k-r)}(z)\restriction_M \\ &= \sigma^{t-(j_k-i_k-r)}(\tau)\restriction_M .\end{aligned}$$

This contradicts the assumption that $\tau_1 \dots \tau_M$ never reoccurs in τ. The case that α_k satisfies (2) of Definition 3.7 is similar (notationally much simpler). The proof follows for non-recurrent τ.

This will lead to another precritical point z' which is responsible for the flip. The fact that the flip occurs between $j_k - i_k + 1$ and $(j_k - i_k + 1) + N_\varepsilon$ implies that there we must have some t such that $\sigma^t(z') = *\tau_1\tau_2\dots$ and such

Next assume that τ is recurrent but not periodic and we have enumerated the return times as in Lemma 3.6. Specifically, let $\{m_i\}_{i\in\mathbb{N}}$ be a sequence of natural numbers chosen such that

$$m_i \leq r_{m_i} < r_{m_i} + 1 < m_{i+1}.$$

Let t be such that $m_t > N_\varepsilon$. Then in this case define $\delta_\varepsilon > 0$ so small that

$$N_{\delta_\varepsilon} > \left(\sum_{q=t}^{t+2N_\varepsilon+1} m_q\right) + 2N_\varepsilon + 1.$$

Let $0 < \delta \leq \delta_\varepsilon$, and let $\{x_i\}_{i\in\mathbb{N}}$ be a δ pseudo-orbit. As in the periodic case of the proof, let $k \in \mathbb{N}$ and consider α_k. First we consider the case that α_k satisfies case (1) of Definition 3.7. Let flip column j_k and associated flip row i_k be as defined. We prove that there are no flips relative to $x_{\alpha_k+i_k}$ between $j_k - i_k$ and $j_k - i_k + N_\varepsilon$ with flip rows between 1 and $\beta_k + N_\varepsilon - i_k$. Lemma 3.2 will imply the proposition.

Suppose that there is a flip column relative to $x_{\alpha_k+i_k}$ in column t_1 with $j_k - i_k < t_1 < j_k - i_k + N_\varepsilon$. Choose t_1 minimally with respect to this property. Then we have that $t_1 - (j_k - i_k) < N_\varepsilon < m_t$. So by Lemma 3.6 there next flip column relative to $x_{\alpha_k+i_k}$ must occur within m_{t+1} of t_1. Letting that flip column be t_2, we have

$$t_2 - t_1 - 1 < m_{t+1}.$$

Therefore,

$$t_2 - t_1 + 1 \leq m_{t+1} + 1 \leq r_{m_{t+1}} + 1 < m_{t+2} \leq r_{m_{t+2}}.$$

So we can iterate the process where in the qth step we get $t_q > t_{q-1}$ is minimal such that t_q is a flip column for $x_{\alpha_k+i_k}$ and

$$t_q - t_{q-1} < m_{t+q-1}.$$

This process must terminate at least by the time we have exhausted the possible flip rows between $x_{\alpha_k+i_k}$ and $x_{\beta_k+N_\varepsilon} = x_{\alpha_k+j_k+N_\varepsilon}$. Let t_r be the last such flip column. Then $r \leq j_k + N_\varepsilon - i_k \leq j_k + N_\varepsilon < 2N_\varepsilon$. Therefore $t_r - t_{r-1} < m_{t+2N_\varepsilon-1}$. This implies that

$$t_r - j_1 = t_r - t_{r-1} + t_{r-1} - \cdots + t_1 - j_1$$

and therefore

$$t_r - j_1 < \sum_{q=t}^{t+2N_\varepsilon-1} m_q.$$

But $N_\delta - j_1 > N_\delta - N_\varepsilon > \left(\sum_{q=t}^{t+2N_\varepsilon} m_q\right) + N_\varepsilon$, so

$$N_\delta - t_r - 1 > m_{t+2N\varepsilon} + N_\varepsilon - 1 > m_r.$$

But in this case we can apply Lemma 3.6 again to construct a flip row t_{r+1} for $x_{\alpha_k+i_k}$ with $t_{r+1} - t_r < m_r$. This contradicts the fact that t_r is the last flip column relative to $x_{\alpha_k+i_k}$. Again, the case that α_k satisfies (2) of Definition 3.7 is similar. The proposition follows. ☐

Proposition 3.9. *Let* $\varepsilon > 0$, *with* $N_\varepsilon \in \mathbb{N}$ *as in Lemma 2.7 and* $\delta_\varepsilon > 0$ *as in Proposition 3.8. If* $\{x_n\}_{n\in\mathbb{N}}$ *is a* δ_ε *pseudo-orbit, where for each* n *we write* $x_n = x_0^n x_1^n \ldots$, *then for each* $n \in \mathbb{N}$

$$x_n{\restriction_{N_\varepsilon}} \simeq x_0^n x_0^{n+1} \ldots x_0^{n+N_\varepsilon}.$$

Proof. Pick $n \in \mathbb{N}$. If there is no $k \in \mathbb{N}$ with $n \leq \beta_k \leq N_\varepsilon$ we have that

$$x_n{\restriction_{N_\varepsilon}} = x_0^n x_0^{n+1} \ldots x_0^{n+N_\varepsilon}$$

and we are done. So assume that there is some minimal β_k between n and N_ε, which represents the first flip column j_k relative to x_i, with flip row i_k. By Proposition 3.8,

$$x_{\beta_k+t}{\restriction_{N_\varepsilon-t}} \approx \sigma^t(*\tau){\restriction_{N_\varepsilon-t}} \tag{4}$$

for all $0 \leq t \leq N_\varepsilon$. Moreover, since j_k and i_k are minimal we have that

$$x_r^n = x_0^{n+r} \tag{5}$$

for $0 \leq r < j_k$, and $x_{j_k+s}^n = x_{j_k-i_k+s+1}^{n+i_k-1}$; in other words that

$$\sigma^{j_k}(x_n){\restriction_{N_\varepsilon-j_k}} = \sigma^{j_k-i_k+1}(x_{n+i_k-1}){\restriction_{N_\varepsilon-j_k}} . \tag{6}$$

Since there is a flip in row i_k relative to x_n,

$$\begin{aligned}\sigma^{j_k-i_k+1}(x_{n+i_k-1})\restriction_{N_\varepsilon-j_k} &\approx *\tau\restriction_{N_\varepsilon-j_k}\\ &\approx x_{\beta_k}\restriction_{N_\varepsilon-j_k}\end{aligned}$$

by (4). The result follows by (5) and (6). □

4. Shadowing and ω-Limit Sets in Dendrites

In this section we prove that for a Λ-acceptable τ, the shift map on the dendrite $\mathcal{D}_\tau$ has shadowing, and we use this fact to prove the main theorem: a closed set $B \subset \mathcal{D}_\tau$ is an ω-limit set of a point if, and only if, B is internally chain transitive (ICT).

Recall the definition of the sequences $\{\alpha_k\}_{k\in\mathbb{N}}$ and $\{\beta_k\}_{k\in\mathbb{N}}$ from Definition 3.7.

Definition 4.1. Let $\varepsilon > 0$, let $N_\varepsilon \in \mathbb{N}$ be given by Lemma 2.7 and $\delta_\varepsilon > 0$ be given by Proposition 3.8. Let $x_1x_2\dots$ be a δ_ε pseudo-orbit (finite or infinite), for each i write $x_i = x_0^i x_1^i \dots$ and define $\hat{z} = \hat{z}_0\hat{z}_1\dots$ by

$$\hat{z}_i = \begin{cases} x_0^i & \text{if } i \neq \beta_k \text{ for any } k \in \mathbb{N},\\ \diamond & \text{if } i = \beta_k \text{ for some } k \in \mathbb{N}.\end{cases}$$

If the pseudo-orbit is finite with last point x_n, let $\hat{z}_{n+i} = x_i^n$ for every $i \geq 0$. The sequence $\hat{z}$ is called a *canonical ε-shadow* for the δ_ε pseudo-orbit $\{x_i\}$.

Theorem 4.2. *Let τ be Λ-acceptable. Then the shift map on the dendrite $\mathcal{D}_\tau$ has shadowing.*

Proof. For a given $\varepsilon > 0$, let N_ε and δ_ε be as given in Definition 4.1, let $\{x_i\}_{i\in\mathbb{N}}$ be a δ_ε pseudo-orbit, and let $\hat{z}$ be the canonical ε-shadow for $\{x_i\}_{i\in\mathbb{N}}$. Let $z \in \{0,1,*\}^\omega$ be such that $z_i = \hat{z}_i$ for all $i \neq \beta_k$. If for all $k \in \mathbb{N}$, $\sigma^{\beta_k}(\hat{z}) \not\approx \tau$ let z_{β_k} be either 0 or 1; if instead there is some least k such that $\sigma^{\beta_k}(\hat{z}) \approx \tau$ then define $z_{\beta_k} = *$ and for all $i > \beta_k$ let $z_i = \tau_{i-\beta_k}$. Defined in this way, it is clear that $z \in \mathcal{D}_\tau$.

We show that $d(\sigma^i(z), x_i) < \varepsilon$ for every $i \in \mathbb{N}$, which by Lemma 2.7 is equivalent to saying that

$$\sigma^i(z)\restriction_{N_\varepsilon} \simeq x_i\restriction_{N_\varepsilon}.$$

Pick $i \in \mathbb{N}$, then by Proposition 3.9

$$x_i\restriction_{N_\varepsilon} \simeq x_0^i x_0^{i+1} \dots x_0^{i+N_\varepsilon}$$

Thus $\sigma^i(z)\restriction_{N_\varepsilon} \simeq x_i\restriction_{N_\varepsilon}$ by the definition of the canonical shadow, and so $d(\sigma^i(z), x_i) < \varepsilon$. □

We now prove our main theorem, which is the following.

Theorem 4.3. *Let* τ *be* Λ*-acceptable. Then* $B \subset \mathcal{D}_\tau$ *is closed and internally chain transitive if, and only if,* $B = \omega(z)$ *for some* $z \in \mathcal{D}_\tau$*.*

Proof. $\mathcal{D}_\tau$ has shadowing by Theorem 4.2. For every $i \in \mathbb{N}$ let δ_i be the constant given by shadowing such that each δ_i pseudo-orbit is $1/2^i$-shadowed.

Suppose that $B \subset \mathcal{D}_\tau$ is closed and internally chain transitive. If B is finite, to be ICT it must be a cycle and is thus an ω-limit set, so assume that B is infinite; then to be ICT it must contain infinitely many non-precritical points.

By compactness of B let $\{x_i\}_{i\in\mathbb{N}}$ be a dense subset of B, then since B is ICT, for every $i \in \mathbb{N}$ there is a δ_i pseudo-orbit $\Gamma_i = \{x_i = x_1^i, \dots, x_{n_i}^i = x_{i+1}\}$, with $n_i > N_{1/2^{i-1}}$ for $N_{1/2^{i-1}}$ given by Lemma 2.7. (If the δ_i pseudo-orbit between x_i and x_{i+1} has length less than $N_{1/2^{i-1}}$ we can add extra δ_i pseudo-orbits from x_{i+1} to itself until the inequality is satisfied.)

For each $i \in \mathbb{N}$, let $z_i = z_0^i z_1^i z_2^i \dots$ be an assignment of the canonical $1/2^i$-shadow for Γ_i as constructed in the proof of Theorem 4.2, except that we do not assign a $*$ to any $\diamond$ in any z_i. Define $z \in \{0,1\}^\omega$ by

$$z = z_0^1 z_1^1 \dots z_{n_1-1}^1 z_0^2 z_1^2 \dots z_{n_2-1}^2 z_0^3 z_1^3 \dots$$

and let $\nu_i = \sum_{j\le i} n_j$. We claim that $z \in \mathcal{D}_\tau$ and that $\omega(z) = B$.

To see that $z \in \mathcal{D}_\tau$, notice first that since B contains infinitely many non-precritical points, elements of $\{z_i\}_{i\in\mathbb{N}}$ will contain arbitrarily long initial segments of non-precritical points, and we deduce that $\sigma^k(z) \neq \tau$ for any $k \in \mathbb{N}$. If $z \notin \mathcal{D}_\tau$, then by the definition of $\mathcal{D}_\tau$ we must have that for some $k \in \mathbb{N}$, $\sigma^k(z)$ only differs from τ in a place where one has a $*$; z has no $*$ so τ must have a $*$ in this place, and as such is periodic. But then by the construction of z, cofinitely many of the z_i would have to be precritical, forcing B to be the critical cycle, a contradiction.

To see that $B \subset \omega(z)$, notice that $\sigma^{\nu_i}(z)\restriction_{N_{1/2^i}} = z_i\restriction_{N_{1/2^i}}$ for every i, so

$$d(\sigma^{\nu_i}(z), x_i) < \frac{1}{2^i}$$

for every $i \in \mathbb{N}$ by the definition of the canonical shadow. For $y \in B$ and $\varepsilon > 0$, pick $j \in \mathbb{N}$ so that $\max\{d(x_j, y), 1/2^j\} < \varepsilon/2$. Then

$$\begin{aligned} d(\sigma^{\nu_j}(z), y) &\le d(\sigma^{\nu_j}(z), x_j) + d(x_j, y) \\ &< \frac{1}{2^j} + d(x_j, y) \\ &< \varepsilon/2 + \varepsilon/2, \end{aligned}$$

so $y \in \omega(z)$.

To see that $\omega(z) \subset B$, notice first that by the definition of the canonical shadow for finite pseudo-orbits, each z_i begins with a length-$N_{1/2^{i-1}}$ portion of x_i and ends with the whole of x_{i+1}, so we immediately see that there is an $N_{1/2^{i-1}}$ overlap between z_i and z_{i+1} for every $i \in \mathbb{N}$.

Suppose for a contradiction that there is some $y \in \omega(z) \setminus B$. Since B is compact there is some $\eta > 0$ such that $y \notin B_\eta(B)$; pick $J \in \mathbb{N}$ such that $1/2^J < \eta$. Since $y \in \omega(z)$, for infinitely many $i > J$ we must have that

$$\sigma^{k_i}(z)\restriction_{N_{1/2^i}} \simeq y\restriction_{N_{1/2^i}}, \tag{7}$$

for appropriate integers k_i, by Lemma 2.7. Since $y \notin B$, $y\restriction_{N_\eta} \not\simeq x\restriction_{N_\eta}$ for all $x \in B$. But then by (7), for infinitely many $i > J$,

$$\sigma^{k_i}(z)\restriction_{N_\eta} \not\simeq x\restriction_{N_\eta}$$

for any $x \in B$, meaning that for infinitely many z_i,

$$\sigma^{j_i}(z_i)\restriction_{N_\eta} \not\simeq x\restriction_{N_\eta}\ .$$

This contradicts the fact that the z_i $1/2^i$-shadow points in B, so $y \in \omega(z)$ and we are done. ☐

Finally, Theorems 2.3 and 4.3 imply the following:

Corollary 4.4. *For any quadratic map $f : J_c \to J_c$ with a dendritic Julia set J_c, $\Lambda \subset J_c$ is closed and internally chain transitive if and only if $\Lambda = \omega(z)$ for some $z \in J_c$.*

REFERENCES

[1] L. Alsedà and N. Fagella. Dynamics on Hubbard trees. *Fund. Math.*, 164(2):115–141, 2000.

[2] S. Baldwin. Continuous itinerary functions and dendrite maps. *Topology Appl.*, 154(16):2889–2938, 2007.

[3] S. Baldwin. Inverse limits of tentlike maps on trees. *Fund. Math.*, 207(3):211–254, 2010.

[4] S. Baldwin. Julia sets and periodic kneading sequences. *J. Fixed Point Theory Appl.*, 7(1):201–222, 2010.

[5] A. Barwell, C. Good, R. Knight, and B. E. Raines. A characterization of ω-limit sets in shift spaces. *Ergodic Theory Dynam. Systems*, 30(1):21–31, 2010.

[6] A. D. Barwell. A characterization of ω-limit sets of piecewise monotone maps of the interval. *Fund. Math.*, 207(2):161–174, 2010.

[7] A. D. Barwell, C. Good, and G. Davies. On the ω-limit sets of tent maps. *Fund. Math.*, (to appear).

[8] A. D. Barwell, C. Good, P. Oprocha, and B. E. Raines. Characterizations of ω-limit sets of topologically hyperbolic spaces. *arXiv*, (1110.3219), 2011.

[9] A. Douady and J. H. Hubbard. *Étude dynamique des polynômes complexes. Partie I*, volume 84 of *Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]*. Université de Paris-Sud, Département de Mathématiques, Orsay, 1984.

[10] M. W. Hirsch, Hal L. Smith, and X.-Q. Zhao. Chain transitivity, attractivity, and strong repellors for semidynamical systems. *J. Dynam. Differential Equations*, 13(1):107–131, 2001.

[11] D. Schleicher. On fibers and local connectivity of Mandelbrot and Multibrot sets. In *Fractal geometry and applications: a jubilee of Benoît Mandelbrot. Part 1*, volume 72 of *Proc. Sympos. Pure Math.*, pages 477–517. Amer. Math. Soc., Providence, RI, 2004.

[12] P. Walters. On the pseudo-orbit tracing property and its relationship to stability. In *The structure of attractors in dynamical systems (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977)*, volume 668 of *Lecture Notes in Math.*, pages 231–244. Springer, Berlin, 1978.

(A. D. Barwell) School of Mathematics, University of Bristol, Howard House, Queens Avenue, Bristol, BS8 1SN, UK – and – School of Mathematics, University of Birmingham, Birmingham, B15 2TT, UK

E-mail address, A. D. Barwell: `A.Barwell@bristol.ac.uk`

(B. E. Raines) Department of Mathematics, Baylor University, Waco, TX 76798–7328,USA

E-mail address, B. E. Raines: `brian_raines@baylor.edu`